

Contains: disodium tetraborate pentahydrate, borax pentahydrate

CAS: 12179-04-3

WARNING - H272: May intensify fire; oxidiser. H315: Causes skin irritation. H319: Causes serious eve irritation. H361: Suspected of damaging fertility or the unborn child. P201: Obtain special instructions before use. P210: Keep away from heat, hot surfaces, open flames, sparks. - No smoking. P221: Take any precaution to avoid mixing with combustibles. P280: Wear face shield, protective gloves. P305+P351+P338: If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P313: If exposed or concerned: Get medical advice/attention.

Manufactured in Spain by: Timac Agro España S.A. Polígono Arazuri-Orcoyen -Calle C, n°32 ORCOYEN (NAVARRA) 31160 ESPAÑA - Tel: + 34 948 324 500

# CORONA<sup>TM</sup>K

Guaranteed analysis	F1550	
Total nitrogen (N): 8% 8% Nitrate nitrogen (N)	Manganese (Mn) water soluble 0.1% 0,1% Chelated Manganese (Mn)	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ): 11% Soluble Potash (K <sub>2</sub> O): 39% Boron (B) water soluble 0.1%	Zinc (Zn).water soluble	
Copper (Cu) water soluble	Monopotassium phosphate, Potassium sulfate, Ammonium sulfate, Boron pentahydrate, Copper EDTA, Iron EDTA, Manganese EDTA, Zinc EDTA.	

### PLEASE READ CAREFULLY BEFORE USING DİRECTİONS FOR USE: KEEP OUT OF REACH OF CHILDREN

Crop	N° of applications	Application stages	Application rate
All row crops including Soybeans, Corn, Wheat, Grains, Peanuts, all potatoes	1-3	Developmental and post bloom	1 – 4 lbs/A
Citrus	1 – 10	Pre bloom , post bloom, summer, fall spray	1 – 10 lbs/A
All fruits and nut crops	1 – 10	Pre bloom, bloom, summer and fall flushes	1 – 10 lbs/A
All vegetables	1 – 10	Development stages ,pre bloom, correct deficiencies	1 – 10 lbs/A
Turfgrasses and Ornamentals	1 – 3	For finish quality As Needed	Turf: 2 – 10 lbs/A Ornamental: 2-3 lbs/A

INDICATIONS DOSAGE AND CONDITIONS OF USE: Add CORONA K first in a half tank water and agitate while mixing. Spray Oils should be added very last to any mix so water solubles can dissolve completely first. Keep agitating until the end of the application. Dilute into at least 15-20 gallons/Acre. It can also be used in ULV Aerial application. Foliar fertilization is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants. Do not spray during periods of frost. In hot conditions spray at the beginning or at the end of the day.

COMPATIBILITY AND STORAGE: CORONA K is compatible with many chemicals. In all cases, test the compatibility prior to treatment. Storage: Keep out of moisture.

WARNING: Some crops may be injured by application of Boron. Net Weight: 13.22 lbs. (6 kg) Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Timac Agro USA, INC. - P.O. Box 888 - Route 724 & I-176 Reading, PA 19607-0888 - USA

CONON-USAG 1/10 - Made III Spain

### Corona N

F 1550 **GUARANTEED ANALYSIS:** Total nitrogen (N) ...... .....21% 8% nitrate Nitrogen (N) 13% ammoniacal Nitrogen (N) Available Phosphate (P2O5)......7% Soluble Potash (K2O) ......7% Magnesium (Mg)......1.2% Sulfur (S)......7.2% Boron(B) .... Water soluble Boron (B) pentahydrate Copper (Cu) ...... 0.1% water soluble chelated Copper (Cu) Iron (Fe) ......0.1% 0.1% chelated Iron (Fe) Manganese (Mn) ......0.1% 0.1% water soluble chelated Manganese (Mn) Molybdenum (Mo)...... 0.01% Zinc (Zn)...... 0.1% chelated Zinc (Zn)

DERIVED FROM: Boron pentahydrate, Monoammonium Phosphate, Sodium Molybdate dihydrate, Ammonium Nitrate, Potassium Nitrate, Ammonium sulfate, Magnesium sulfate, Copper EDTA chelate, Iron EDTA chelate, Manganese EDTA chelate, Zinc EDTA chelate.

WARNING: Some other crops may be injured by application of Boron (B). This product contains molybdenum (Mo). Crops high in molybdenum can be toxic to grazing animals. Use only according to manufacturer's directions.

#### DIRECTIONS FOR USE:

- Use in accordance with recommendation from a qualified individual or institution, or according to an approved nutrient management plan.
- Fill tank half way, add product and agitate until fully dissolved. Spray oils should be added last.
- For ground application dilute into at least 15 20 gal/ac. For ULV aerial application use 3 5 gal/ac.
- Keep agitating until the end of the application.

#### PRODUCT USEAGE:

Row Crops: 1-3 applications/year at 1-5 lbs /ac

- Specialty Crops: 1-10 applications/year 2-10 lbs ac
- Foliar fertilization is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required.
- Do not spray during periods of frost. In hot conditions spray at the beginning or at the end of the day.

COMPATIBILITY AND STORAGE: The product is compatible with many chemicals. In all cases, test the compatibility prior to treatment. Storage: Keep out of moisture.

NET WEIGHT: 13.22 lbs. (6 kg)


#### Timac Agro USA - PO Box 888 - Route 724 & 1176 Reading, PA 19607-0888 USA

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.



H319 - Causes serious eye irritation. H360 - May damage fertility or the unborn child (oral). P201 - Obtain special instructions before use. P280 - Wear face shield, protective clothing, protective gloves. P308+P313 - If exposed or concerned: Get medical advice/attention. P337+P313 - If eye irritation persists: Get medical advice/attention. P501 - Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.

Manufactured by: Timac Agro España S.A. - Polígono Arazuri-Orcoyen -Calle C, n°32 -ORCOYEN (NAVARRA), 31160 ESPAÑA



Contains: manganese sulfate, zinc sulfate

CAS: 7785-87-7 CAS: 7733-02-0

DANGER - H318: Causes serious eve damage. H360: May damage fertility or the unborn child, H372: Causes damage to organs through prolonged or repeated exposure (Inhalation, oral). P201: Obtain special instructions before use. P260: Do not breathe dust. P280: Wear face shield, protective clothing, protective gloves. P305+P351+P338: If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P313: If exposed or concerned: Get medical advice/attention. P310: Immediately call a doctor, a POISON CENTER, P501: Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.

Manufactured in Spain by: Timac Agro España S.A. Polígono Arazuri-Orcoven - Calle C, n°32 ORCOYEN (NAVARRA) 31160 ESPAÑA - Tel: + 34 948 324 500

## CORONA MASTER<sup>TM</sup>

8-27-0

Guaranteed analysis Total nitrogen (N): 8% 6% Ammoniacal nitrogen (N) 2% Urea nitrogen (N) Available Phosphate (P<sub>2</sub>O<sub>5</sub>): 27% Magnesium (Mg): 0.9 % Sulfur (S): 8% 0.5% Water Soluble Boron (B) Copper (Cu) ...... 1%

1% Water Soluble Copper (Cu)

DİRECTİONS FOR USE:

Manganese (Mn)...... 6% 6% Water Soluble Manganese (Mn) Molybdenum (Mo)...... 0.1% 0.1% Water Soluble Molybdenum (Mo) 5% Water Soluble Zinc (Zn) DERIVED FROM Monoammonium phosphate, Manganese sulfate, Boron pentahydrate, Zinc sulfate, Magnesium sulfate, Urea, Ammonium sulfate, Copper sulfate, Sodium molybdate.

#### PLEASE READ CAREFULLY BEFORE USING KEEP OUT OF REACH OF CHILDREN

Crop	N° of applications	Application stages	Application rate
All row crops including Soybeans, Corn, Wheat, Grains, Peanuts, all potatoes	1 – 3	Developmental and post bloom	2 – 4 lb/A
Citrus	1 – 10	Pre bloom , post bloom, summer, fall spray	1 – 10 lb/A
All fruits and nut crops	1 – 10	Pre bloom, bloom, summer and fall flushes	1 – 10 lb/A
All vegetables	1 – 10	Development stages ,pre bloom, correct deficiencies	1 – 10 lb/A
Turfgrasses and Ornamentals	1-3	For plant establishment	2 – 3 lb/A

INDICATIONS DOSAGE AND CONDITIONS OF USE: Add CORONA Master first in a half tank water and agitate while mixing. Spray Oils should be added very last to any mix so water solubles can dissolve completely first. Keep agitating until the end of the application. Dilute into at least 15-20 gallons/Acre. Foliar fertilization is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants. Do not spray during periods of frost. In hot conditions spray at the beginning or at the end of the day.

COMPATIBILITY AND STORAGE: CORONA Master is compatible with many chemicals. In all cases, test the compatibility prior to treatment. Storage: Keep out of moisture.

Warning: Some crops may be injured by application of Boron. This product contains molybdenum (Mo). Crops high in molybdenum can be toxic to ruminant animals. Use only according to manufacturer's directions.

Net Weight: 13.22 lbs (6 kg)

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc., In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Timac Agro USA, INC. - P.O. Box 888 - Route 724 & I-176 Reading, PA 19607-0888 - USA