



Maxifruit®

264 Gal.

MAXIFRUIT® 3-7-7

	1417 (7411 1
GUARANTEED ANALYSIS	
Total Nitrogen (N)	3%
3% Urea Nitrogen (N)	
Available Phosphate (P2O5)	7%
Soluble Potash (K2O)	7%
Manganese (Mn)	0.05%
0.05% Water Soluble Manganese (Mn)	
• Zinc (Zn)	0.1%
0.1% Water Soluble Zinc (Zn)	

F 1550

NON PLANT FOOD INGREDIENTS:

MAXIFRUIT® complex10 %

DERIVED FROM: Urea, Phosphoric acid, Potassium hydroxide, Manganese EDTA chelate, Zinc EDTA chelate.

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN - RECIRCULATE WELL BEFORE USE

DIRECTIONS FOR USE:

Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Fill tank half full with clean water then add MAXIFRUIT® and continue to agitate while mixing. Keep agitating until the end of the application. Use the dilution rate in the table below.

• MAXIFRUIT® is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

CROP	N° OF APP.	APPLICATION STAGES	APP. RATE (pts/Acre)	DILUTION (Gal/Acre)
Artichokes	3	Pre-flower induction (before head formation)	2 – 2.5	100
		Shaped head		
		Head growth		
Citrus	1 – 2	Petals falling	1.5 – 2	40 – 100
		8 days after		
Cucurbits	2	10 days before flowering	1.5 – 2	
(cucumbers, zucchini,				40 – 100
melons, watermelons)		Flowering		
Kiwi	2-3	Before flowering	1.5 – 2	40 – 100
		Petals falling		
		15 days after		
Olives	1 – 2	Before flowering	2 – 2.5	90 – 100
		Fruit setting		
Pears and Apples	2 – 3	Before flowering	1.5 – 2	40 – 100
		Petals falling		
		15 days after		
Stone fruits	2-3	Petals falling	1.5 – 2	40 – 100
(peach, apricots, cherries)		15 days after		
Table grapes	2-3	Before flowering (grapes 2"-4")	1.5 – 2	40 – 100
		Fruit setting		
		Fruits 10 mesh		
Tomatoes	3	Before flowering	2 – 2.5	100
		After fruit setting		
		Fruit growing		
Vinyard Grapes	2	Before flowering (grapes 2"-4")	2 – 2.5	40 – 100
		Equit cotting	15 2	

NET VOLUME: 264 Gal (1000L) - NET WEIGHT: 2645 lb (1205 kg) Timac Agro USA, Inc. marrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express of implied warranty except as stated above.

Timac Agro USA, INC. – P.O. Box 888 – Route 724 & 1-176 Reading, PA 19607-0888 – USA

MANUFACTURED BY TIMAC AGRO USA -P.O. Box 888 - Route 724 & I-176 Reading, PA 19607 -0888 - USA

