

FERTIACTYL[®] CORN PLUS

10-0-5

FERTIACTYL IS A LIQUID NUTRITIONAL DESIGNED TO
HELP INCREASE ROOT GROWTH, CROP STAND AND
LESSEN EARLY SEASON STRESS.

F1550

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 10%

10% UREA NITROGEN (N)

SOLUBLE POTASH (K₂O) 5%

ZINC (Zn) 5%

DERIVED FROM: UREA, POTASSIUM HYDROXIDE AND
AMMONIATED ZN.

NON PLANT FOOD INGREDIENTS: HUMIC ACID
(10%), FULVIC ACID (5%), SEAWEED EXTRACT
(20%), OTHER INERT INGREDIENTS (45%).

**KEEP OUT OF REACH OF CHILDREN -
READ ENTIRE LABEL BEFORE USING THIS
PRODUCT - SHAKE OR RECIRCULATE
BEFORE USE**

DIRECTIONS FOR USE: FIELD, ROW, VEGETABLE,
ORCHARD AND VINE CROPS: APPLY 1-4 QUARTS PER ACRE
PER APPLICATION THROUGHOUT THE GROWING SEASON. MORE
FREQUENT APPLICATIONS AT 1 QT./AC PER ACRE MAY BE NEEDED
TO CORRECT DEFICIENCIES OR ON CROPS THAT UNDER HEAVY
WEATHER STRESS. DO NOT SPRAY DURING PERIODS OF FROST, IN
HOT CONDITIONS IT IS BETTER TO SPRAY AT THE BEGINNING OR
AT THE END OF THE DAY.

H315: CAUSES SKIN IRRITATION.

H319: CAUSES SERIOUS EYE IRRITATION.

H335: MAY CAUSE RESPIRATORY IRRITATION.

P271: USE ONLY OUTDOORS OR IN A WELL-VENTILATED AREA.

P280: WEAR PROTECTIVE GLOVES/PROTECTIVE CLOTHING/EYE PRO-
TECTION/FACE PROTECTION.

P302 + P352: IF ON SKIN: WASH WITH PLENTY OF SOAP AND
WATER.

P304 + P340: IF INHALED: REMOVE PERSON TO FRESH AIR AND
KEEP COMFORTABLE FOR BREATHING.

P305 + P351 + P338 + P310: IF IN EYES: RINSE CAUTIOUSLY WITH
WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT
AND EASY TO DO. CONTINUE RINSING. IMMEDIATELY CALL A DOCTOR, A
POISON CENTER.

P337 + P313: IF EYE IRRITATION PERSISTS: GET MEDICAL ADVICE/
ATTENTION.



WARNING

FERTIACTYL CORN PLUS

CROPS	APPLICATION STAGE	APPLICATION RATE 1-12 APPLICATIONS
ORCHARDS AND VINEYARDS INCLUDING BUT NOT LIMITED TO: APPLES, PEARS, PEACHES, NECTARINES, GRAPES, (JUICE, TABLE, WINE), CITRUS	FOLIAR: BUDBREAK// HARVEST GROUND APPLICATION	1-3 QT./AC FOLIAR 2-4 QT./ACRE SOIL APPLIED AS OFTEN AS WEEKLY
ROOT AND VEGETABLES, INCLUDING BUT NOT LIMITED TO: SUGAR BEETS, POTATOES, CARROTS, ONIONS, AND OTHER ROOT CROPS, TOMATOES, PEPPERS, EGGPLANTS, CUCURBITS, PUMPKINS, SQUASHES, CARROTS, LEAFY GREENS, COLD CROPS (CABBAGE, BRUSSEL SPROUTS, BROCCOLI)	WITH LIQUID SIDE-DRESS OR THROUGH FERTIGATION FOLIAR: EMERGENCE TO V6	1-2 QTS./AC FOLIAR 2-4 QTS./AC SOIL APPLIED AS OFTEN AS BI-WEEKLY
FIELD AND ROW CROPS, INCLUDING BUT NOT LIMITED TO: CORN, WHEAT, BARLEY, RYE, OATS, SORGHUM, OTHER CEREAL GRAINS, SUNFLOW- ER, CANOLA, SOYBEAN, LENTILS, DRY BEANS, FIELD PEAS, PEANUTS, COTTON, RICE	WITH LIQUID SIDE-DRESS, STARTER FERTILIZER, 2x2, OR IN-FURROW FOLIAR: EMERGENCE TO V6	1-2 QTS./AC FOLIAR 0.5-2 QT./AC SOIL APPLIED
ALFALFA, CLOVER, AND GRASS CROPS	FOLIAR APPLY AT SPRING GREEN UP AND AGAIN 3-7 DAYS AFTER EACH CUTTING	1-2 QT./AC FOLIAR 1-4 QT./ACRE SOIL APPLIED

THESE RECOMMENDATIONS ARE GUIDELINES ONLY. OPTIMUM RATE OF APPLICATION WILL VARY DEPENDING ON SOIL PROPERTIES
(SUCH AS pH, ORGANIC MATTER CONTENT, TEXTURE), WEATHER CONDITIONS, TIME OF YEAR, PLANT SPECIES AND GENERAL HEALTH.
FOR BEST RESULTS, FOLLOW SOIL AND TISSUE TEST RECOMMENDATIONS.

COMPATIBILITY, TANK MIXING AND STORAGE: CHECK MIXTURE COMPATIBILITY USING A JAR TEST PRIOR TO
APPLICATION. FILL MIX TANK WITH HALF OF REQUIRED WATER AND BEGIN AGITATION. ADD PRODUCTS IN THIS ORDER, MIXING
THOROUGHLY AFTER EACH ADDITION: ADJUVANTS, PESTICIDES, THEN FERTILIZERS. FILL TANK WITH REMAINDER OF WATER AND
CONTINUE AGITATION UNTIL SOLUTION IS COMPLETELY MIXED. MANUFACTURER RESPONSIBILITY IS LIMITED TO THE GUARANTEE OF THE
PRODUCT'S COMPOSITION AS PER LABEL IN CERTIFIED PACKAGING. IN ALL CASES, TEST THE COMPATIBILITY PRIOR TO TREATMENT.
STORE IN A COOL AND DRY PLACE, AVOID STORING IN SUNLIGHT OR HIGH HEAT ENVIRONMENTS. DO NOT STORE UNDER 32°F
(0°C).

NOTICE OF WARRANTY: TIMAC AGRO USA, INC. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL
DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR PURPOSES STATED ON THE LABEL WHEN USED IN ACCORDANCE
WITH DIRECTIONS UNDER NORMAL USE CONDITIONS. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES
MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE
OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF TIMAC AGRO USA, INC. IN NO CASE SHALL TIMAC AGRO
USA, INC. BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR SHIPPING
OF THIS PRODUCT. ALL SUCH RISK SHALL BE ASSUMED BY THE BUYER. TIMAC AGRO USA, INC. MAKES NO WARRANTIES OF
MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED
ABOVE.


WEIGHT PER GALLON

10.95 lbs.

NET CONTENT

250 Gal | 2,737 lbs. (1241 kg)

MANUFACTURED BY TIMAC AGRO USA
P.O. Box 888 - ROUTE 724 & I-176 READING, PA
19607 -0888 - USA



FERTIACTYL®



GZ



264 Gal.

FERTIACTYL® GZ 13-0-5

GUARANTEED ANALYSIS

- Total Nitrogen (N) 13%
13 % Urea Nitrogen
- Soluble Potash (K₂O) 5%
DERIVED FROM : Urea, Potassium Hydroxide

F 1550

NON PLANT FOOD INGREDIENTS:

- Humic acid 10 %
- Fulvic acid 5 %
- Seaweed extract 20 %

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN - RECIRCULATE BEFORE USE

DIRECTIONS FOR USE:

- Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Add FERTIACTYL® GZ first in a half full clean sprayer and agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 Gal/Acre.
- FERTIACTYL® GZ is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

CROP	N° OF APP.	APPLICATION STAGES	APP. RATE (pts/Acre)	ADVICE
Corn, Soybean & Alfalfa	1 - 4	At planting	2 - 4	Broadcast app.
			1.5 - 4	In row app.
		Early growth	2 - 4	Broadcast app.
			1.5 - 4	Banded app.
Cotton	2	At planting	2 - 4	• Apply in furrows. • 4 pts/Acre gives best results. • It can be added to liquid starter fert. with glyphosate foliar app. Total should be 4 pts/Acre.
		With herbicide application		
General Vegetables	1 - 3	At planting	0.25 - 1% transplant solution	Must have mechanical or by pass tank agitation in transplant water tank.
		At transplant to 7 days after planting	2 - 4	May be mixed with compatible fertilizers (not calcium nitrate) or injected in clean water (drip irrigation). If not available, use it to water plants (same rate), wetting leaves until the solution reaches the soil.
		With 2nd foliar spray (7 - 14 days after 1st application)		
Wheat and Small Grains	1 - 2	Planting to FS 3	1 - 3	• Tank mixes well with most pesticides and liquid fertilizers. • Enhances tillering.

COMPATIBILITY AND STORAGE:

- FERTIACTYL® GZ is compatible with most liquid fertilizers and a wide range of agrochemicals. Do not mix with low pH products or calcium nitrate. In all cases, test the compatibility prior to treatment. Manufacturer's responsibility is limited to the guarantee of product's composition, as per label in certificate packaging.
- Do not store under 32°F (0°C).

Volume: 264 Gallons (1000 liters) - Net weight: 2667 lb (1210 kg)

TIMAC AGRO USA, INC. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA, INC. In no case shall TIMAC AGRO USA, INC. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. TIMAC AGRO USA, INC. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY TIMAC AGRO USA -
P.O. Box 888 - Route 724 & I-176 Reading,
PA 19607 - 0888 - USA



FERTIACTYL®



Record

264 Gal.

FERTIACTYL® Record 0-0-30

GUARANTEED ANALYSIS

F 1550

- Soluble Potash (K₂O)..... 30%
- Sulfur (S) 12%

DERIVED FROM: Potassium thiosulfate and Tripotassium citrate.

**PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN
RECIRCULATE WELL BEFORE USE**

DIRECTIONS FOR USE:

- FERTIACTYL® Record may be used in application to the soil, near the root system. Fertigation system or other ways of spreading the product can be used, like overhead irrigation.
- Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Fill tank half full with clean water then add FERTIACTYL® Record and continue to agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 Gal/Acre.
- FERTIACTYL® Record is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

CROP	N° OF APPLICATIONS	APPLICATION STAGES	APPLICATION RATE (pts/Acre)
Citrus & Fruits	1 – 2	Fertigation 9 – 14 days prior to harvest	4
Cucumbers, Melons, Peppers, Potatoes, Strawberries & Tomatoes	1 – 3	From 1 st fruits to early maturation	4 – 8

COMPATIBILITY AND STORAGE:

- FERTIACTYL® Corn is compatible with many chemicals and liquid fertilizers except acid-pH products, calcium nitrate and mineral oil based products. In all cases, test the compatibility prior to treatment.
- Do not store under 32°F (0°C).

NET VOLUME: 264 Gal (1000L) - NET WEIGHT: 3461.3 lb (1570 kg)

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY TIMAC AGRO USA -
P.O. Box 888 - Route 724 & I-176
Reading, PA 19607 - 0888 - USA



FERTIACTYL® RhiZn

10-0-5

FERTIACTYL IS A LIQUID NUTRITIONAL DESIGNED TO
HELP INCREASE ROOT GROWTH, CROP STAND AND
LESSEN EARLY SEASON STRESS.

F1550

GUARANTEED ANALYSIS

TOTAL NITROGEN (N) 10%

10% UREA NITROGEN (N)

SOLUBLE POTASH (K₂O) 5%

ZINC (Zn) 2%

DERIVED FROM: UREA, POTASSIUM HYDROXIDE AND
Zn EDTA.

NON PLANT FOOD INGREDIENTS: HUMIC ACID
(10%), FULVIC ACID (5%), SEAWEED EXTRACT
(20%), OTHER INERT INGREDIENTS (45%).

**KEEP OUT OF REACH OF CHILDREN -
READ ENTIRE LABEL BEFORE USING THIS
PRODUCT - SHAKE OR RECIRCULATE
BEFORE USE**

DIRECTIONS FOR USE: FIELD, ROW, VEGETABLE,
ORCHARD AND VINE CROPS: APPLY 1-4 QUARTS PER ACRE
PER APPLICATION THROUGHOUT THE GROWING SEASON. MORE
FREQUENT APPLICATIONS AT 1 QT/AC PER ACRE MAY BE NEEDED
TO CORRECT DEFICIENCIES OR ON CROPS THAT UNDER HEAVY
WEATHER STRESS. DO NOT SPRAY DURING PERIODS OF FROST, IN
HOT CONDITIONS IT IS BETTER TO SPRAY AT THE BEGINNING OR
AT THE END OF THE DAY.

H315: CAUSES SKIN IRRITATION.

H319: CAUSES SERIOUS EYE IRRITATION.

H335: MAY CAUSE RESPIRATORY IRRITATION.

P271: USE ONLY OUTDOORS OR IN A WELL-VENTILATED AREA.

P280: WEAR PROTECTIVE GLOVES/PROTECTIVE CLOTHING/EYE PRO-
TECTION/FACE PROTECTION.

P302 + P352: IF ON SKIN: WASH WITH PLENTY OF SOAP AND
WATER.

P304 + P340: IF INHALED: REMOVE PERSON TO FRESH AIR AND
KEEP COMFORTABLE FOR BREATHING.

P305 + P351 + P338 + P310: IF IN EYES: RINSE CAUTIOUSLY WITH
WATER FOR SEVERAL MINUTES. REMOVE CONTACT LENSES, IF PRESENT
AND EASY TO DO. CONTINUE RINSING. IMMEDIATELY CALL A DOCTOR, A
POISON CENTER.

P337 + P313: IF EYE IRRITATION PERSISTS: GET MEDICAL ADVICE/
ATTENTION.



WARNING

FERTIACTYL RhiZn

CROPS	APPLICATION STAGE	APPLICATION RATE 1-12 APPLICATIONS
ORCHARDS AND VINEYARDS INCLUDING BUT NOT LIMITED TO: APPLES, PEARS, PEACHES, NECTARINES, GRAPES, (JUICE, TABLE, WINE), CITRUS	FOLIAR: BUDBREAK// HARVEST GROUND APPLICATION	1-3 QT/AC FOLIAR 2-4 QT/ACRE SOIL APPLIED AS OFTEN AS WEEKLY
ROOT AND VEGETABLES, INCLUDING BUT NOT LIMITED TO: SUGAR BEETS, POTATOES, CARROTS, ONIONS, AND OTHER ROOT CROPS, TOMATOES, PEPPERS, EGGPLANTS, CUCURBITS, PUMPKINS, SQUASHES, CARROTS, LEAFY GREENS, COLD CROPS (CABBAGE, BRUSSEL SPROUTS, BROCCOLI)	WITH LIQUID SIDE-DRESS OR THROUGH FERTIGATION FOLIAR: EMERGENCE TO V6	1-2 QTS/AC FOLIAR 2-4 QTS/AC SOIL APPLIED AS OFTEN AS BI-WEEKLY
FIELD AND ROW CROPS, INCLUDING BUT NOT LIMITED TO: CORN, WHEAT, BARLEY, RYE, OATS, SORGHUM, OTHER CEREAL GRAINS, SUNFLOW- ER, CANOLA, SOYBEAN, LENTILS, DRY BEANS, FIELD PEAS, PEANUTS, COTTON, RICE	WITH LIQUID SIDE-DRESS, STARTER FERTILIZER, 2x2, OR IN-FURROW FOLIAR: EMERGENCE TO V6	1-2 QTS/AC FOLIAR 0.5-2 QT/AC SOIL APPLIED
ALFALFA, CLOVER, AND GRASS CROPS	FOLIAR APPLY AT SPRING GREEN UP AND AGAIN 3-7 DAYS AFTER EACH CUTTING	1-2 QT/AC FOLIAR 1-4 QT/ACRE SOIL APPLIED

THESE RECOMMENDATIONS ARE GUIDELINES ONLY. OPTIMUM RATE OF APPLICATION WILL VARY DEPENDING ON SOIL PROPERTIES
(SUCH AS pH, ORGANIC MATTER CONTENT, TEXTURE), WEATHER CONDITIONS, TIME OF YEAR, PLANT SPECIES AND GENERAL HEALTH.
FOR BEST RESULTS, FOLLOW SOIL AND TISSUE TEST RECOMMENDATIONS.

COMPATIBILITY, TANK MIXING AND STORAGE: CHECK MIXTURE COMPATIBILITY USING A JAR TEST PRIOR TO
APPLICATION. FILL MIX TANK WITH HALF OF REQUIRED WATER AND BEGIN AGITATION. ADD PRODUCTS IN THIS ORDER, MIXING
THOROUGHLY AFTER EACH ADDITION: ADJUVANTS, PESTICIDES, THEN FERTILIZERS. FILL TANK WITH REMAINDER OF WATER AND
CONTINUE AGITATION UNTIL SOLUTION IS COMPLETELY MIXED. MANUFACTURER RESPONSIBILITY IS LIMITED TO THE GUARANTEE OF THE
PRODUCT'S COMPOSITION AS PER LABEL IN CERTIFIED PACKAGING. IN ALL CASES, TEST THE COMPATIBILITY PRIOR TO TREATMENT.
STORE IN A COOL AND DRY PLACE, AVOID STORING IN SUNLIGHT OR HIGH HEAT ENVIRONMENTS. DO NOT STORE UNDER 32°F
(0°C).

NOTICE OF WARRANTY: TIMAC AGRO USA, INC. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL
DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR PURPOSES STATED ON THE LABEL WHEN USED IN ACCORDANCE
WITH DIRECTIONS UNDER NORMAL USE CONDITIONS. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES
MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE
OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF TIMAC AGRO USA, INC. IN NO CASE SHALL TIMAC AGRO
USA, INC. BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR SHIPPING
OF THIS PRODUCT. ALL SUCH RISK SHALL BE ASSUMED BY THE BUYER. TIMAC AGRO USA, INC. MAKES NO WARRANTIES OF
MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED
ABOVE.

WEIGHT PER GALLON

10.95 lbs.

NET CONTENT

250 Gal | 2,737 lbs. (1241 kg)

MANUFACTURED BY TIMAC AGRO USA
P.O. Box 888 - ROUTE 724 & I-176 READING, PA
19607 -0888 - USA

